

Preparing for GOES-R and JPSS: New Tropical Cyclone Tools Based on 30 Years of Continuous GOES IR Imagery



John A. Knaff¹, Robert T. DeMaria², Scott P. Longmore², Debra A. Molenar¹, Kate D. Musgrave²

¹NOAA Center for Satellite Applications and Research/RAMMB, Fort Collins, Colorado

²The Cooperative Institute for Research in the Atmosphere, Fort Collins, Colorado


Scientific Premise: As JPSS/SNPP and GOES-R platforms become available, the focus of the satellite community's funding, research, and product development often gravitates toward their new capabilities. This is sometimes justified, especially when the information content of older datasets has been exhausted. However this is sometimes not the case.

Here we argue that tropical cyclone (TC) research and product development is a case where there is untapped potential in the information contained in older satellite data. The reasons are many. The observation of tropical cyclones has dramatically improved in the last 20 years. Some significant observational improvements are based on satellite platforms (e.g., scatterometry, microwave imagery, water vapor imagery) while others (e.g., aircraft reconnaissance instrumentation, improved dropwindsondes) are not. Statistical techniques, data storage, data availability, and computational speed have also all improved. In sum, the availability of improved observations, analysis techniques, computing, and physical understanding of TCs has improved so much that new capabilities can be developed using older geostationary and polar data.

Data Available for TC Application Development

IR Imagery

- CIRA/RAMMB TC IR Image Archive (1990-pres)
- HURSAT (1978-2009)




TC Information

- ATCF (Best tracks w/ wind radii, fixes, guidance, real time)
- HURDAT2 (Best tracks w/ wind radii)
- IBTrACS (all agencies)


Aircraft Reconnaissance

- Air Force/NOAA (10 & 1 Hz; 1995-pres)
- HDOBS (real-time access)
- Dropwindsondes (real-time access)




Other Satellite

- Passive Microwave Imagery/Sounder
- Scatterometry
- CloudSat




Ocean Analyses

- Navy Coupled Ocean Data Assimilation (NCODA)
- NESDIS operational products



Model analysis/forecasts (HWRF/GFS)

- Fields
- Synthetic Imagery



Potential Forecast/Diagnostic Applications

Surface wind structure

- TC size (vortex size as inferred from outer winds)
- Diagnoses of the 2-D wind field using imagery and intensity
- Relating TC size and intensity to wind radii
- Statistical-dynamical wind radii prediction

Eye/radius of maximum winds

- Objective eye detection and/or probability of existence
- Forecasting eye development/eye diameter

Intensity

- Rapid Intensification Forecasts (probabilistic)
- Statistical Dynamical Forecasts (deterministic)


Example Applications Developed with These Data

IR-based TC Size Estimates (Knaff et al. 2014)

Based on the 1D (azimuthally average) Principle components and latitude (no intensity)

$$V500 = 2.488 + 11.478 \sin |\varphi| - 1.350 * PC1 + 0.912 * PC2 + 0.319 * PC3$$

$$R5 = (R5 + (V500 - V500c) * 500 / (V500c - V1000c))$$



Additional Applications Derived from TC Size estimates

Application	Use
1. Wind Radii given intensity, motion vector and TC size	Can be applied to intensity fix(es) (e.g. Dvorak) Provides a distribution for Monte-Carlo/stochastically generated storms for climate risk modeling
2. TC size as the basis for statistical-dynamical size/wind radii forecasts	Can potentially provide wind radii estimates consistent with SHIPS/LGEM
3. TC size can be used to scale observational data	Any application requiring size normalization for average quantities (e.g., rainfall, core/rainband separation, lightning)
4. Model diagnosis	Can quantify sizes from synthetic imagery, assess TC growth etc.


2-D wind field (Knaff et al. 2015)

Based on intensity estimate, 2-D storm relative IR principle components

Trained on aircraft-based flight-level data decomposed into azimuthal wavenumbers 0-1 and their phases

Produces an estimate of the 2-D wind field


Uses a method called single field principle component analysis that relates wind amplitudes and phases to the IR principle components.



Dependent mean wind field (contours) and absolute errors (MAE, shaded) stratified by flight-level analysis winds with cases with winds greater than 96 kt (left), winds 64 to 95 kt (middle) and winds 34 to 64 kt (right). Note motion is toward the top of the page. The magenta contour on each panel shows where the average RMW of flight-level winds is located in each composite. The scale is the same for all figures.


Additional Applications Derived from 2-D Wind Fields

Application	Use
1. NESDIS Multi-Platform Tropical Cyclone Surface Wind Analysis	Improve the inner core flight-level wind estimates
2. Vortex Data assimilation	Can be used to specify the wind distribution including RMW
3. Input for aircraft-based analyses.	First guess



Eye Detection/Eye Size/Radius of Maximum Wind

- Dvorak fixes (ATCF) are used to develop eye no-eye training set (n=2677) based on six-hourly Atlantic and East Pacific tropical cyclones.
- Machine learning algorithms used to train an objective eye detection scheme (under JPSS funding, see Robert DeMaria's poster)
- Preliminary results suggest that this scheme can detect 75% of the eye cases with false detections of ~20% when using a 50 knot intensity threshold



Next Steps

- Refine algorithm with different intensity thresholds/data subsets
- Develop probabilistic eye/no-eye algorithm
- Run the algorithm on all images in the CIRA/RAMMB TC archive (~500,000 unique times)

Additional Applications Eye Detection/Eye Size/Radius of Maximum winds	
Application	Use
1. Intensity forecasting	The probability of eye formation provides vital information as to the short-term intensification rate
2. Climatology of eye size, eye formation	Detailed climatology of eye size and eye formation is of general interest. Eye diameter is related to RMW
3. Eye anticipation, eye size product	This would aid short-term forecasts and forecasters
4. Model diagnosis	Methods could compare model eye formation to climatological aspects of eye formation (2)

Summary

The increased availability of improved observations, new analysis techniques, computing efficiency, and physical understanding of TCs has improved so much that new capabilities can be developed using older geostationary and polar data. Such capabilities can easily be transitioned to future operational and analysis techniques that make use of JPSS and GOES-R imagery and products.

Work to develop new tropical cyclone capabilities using legacy GOES and POES imagery will continue at CIRA/RAMMB and is the basis for much of our GOES-R product development effort.

References:

- Knaff, J.A., S.P. Longmore, D.A. Molenar, 2014: An Objective Satellite-Based Tropical Cyclone Size Climatology. *J. Climate*, **27**, 455-476.
- Knaff, J.A., S.P. Longmore, R.T. DeMaria, D.A. Molenar, 2015: Improved tropical cyclone flight-level wind estimates using routine infrared satellite reconnaissance. Accepted to *J. Appl. Meteor. Climate*.
- DeMaria, R.T., G. Chirokova, J.A. Knaff, and J. F. Dostalek 2015: Machine Learning Algorithms for Tropical Cyclone Center Fixing and Eye Detection. 20th Conference on Satellite Meteorology and Oceanography.

Acknowledgments: This work was supported under the Hurricane Forecast Improvement Program (HFIP) and the GOES-R/JPSS Risk Reduction Program.